

UNCLASSIFIED

---

AD 292 752

*Reproduced  
by the*

ARMED SERVICES TECHNICAL INFORMATION AGENCY  
ARLINGTON HALL STATION  
ARLINGTON 12, VIRGINIA



---

UNCLASSIFIED

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

MEMORANDUM  
RM-3083-PR  
APRIL 1962

292 752

SOME QUESTIONS CONCERNING  
DIFFERENCE APPROXIMATIONS  
TO PARTIAL DIFFERENTIAL EQUATIONS

Richard Bellman

AD No. 32752  
REPRODUCIBLE COPY

ASTIA  
JAN 9 1963  
TISIA A

PREPARED FOR:  
UNITED STATES AIR FORCE PROJECT RAND

#110

The RAND Corporation  
SANTA MONICA • CALIFORNIA

MEMORANDUM

RM-3083-PR

APRIL 1962

SOME QUESTIONS CONCERNING  
DIFFERENCE APPROXIMATIONS  
TO PARTIAL DIFFERENTIAL EQUATIONS

Richard Bellman

This research is sponsored by the United States Air Force under Project RAND — Contract No. AF 49(638)-700 — monitored by the Directorate of Development Planning, Deputy Chief of Staff, Research and Technology, Hq USAF. Views or conclusions contained in this Memorandum should not be interpreted as representing the official opinion or policy of the United States Air Force. Permission to quote from or reproduce portions of this Memorandum must be obtained from The RAND Corporation.

---

*The* **RAND** *Corporation*

1700 MAIN ST • SANTA MONICA • CALIFORNIA

---

PREFACE

Part of the Project RAND research program consists of basic supporting studies. One aspect of this concerns the solution of the partial differential equations of applied mathematics on automatic digital computing machines through approximations by difference equations.

In the present Memorandum the author investigates the determination of discrete approximations that manifestly possess properties, such as boundedness, positivity, and so on, of the given differential operator.

### SUMMARY

The study of finite-difference approximations to partial differential equations is well advanced, and the increasing use and significance of digital computers guarantees that this study will continue to be actively pursued. One question, however, that does not seem to have been discussed to any extent is that of finding discrete approximations that manifestly exhibit the boundedness, positivity, and so on, of the original continuous operator.

Thus, for example, the recurrence relation

$$u(t + \Delta) = (1 - a\Delta)u(t), \quad u(0) = c,$$

yields the nonnegativity we associate with the solution of

$$u' = -au, \quad u(0) = c,$$

provided that  $a\Delta < 1$ . On the other hand, the relation

$$u(x, t + \Delta^2) = \frac{u(x + \Delta\sqrt{2}, t) + u(x - \Delta\sqrt{2}, t)}{2},$$

$t = 0, \Delta^2, \dots$ , shows very clearly the nonnegativity associated with the solution of

$$u_t = u_{xx}, \quad u(x, 0) = g(x).$$

Approximations of this type are very useful computationally because of their stability properties. In this Memorandum, the author discusses briefly the problem of obtaining approximations of arbitrarily high degree with the same property, and the related problem for

$$u_t = uu_x + g(u).$$

CONTENTS

PREFACE . . . . .	.111
SUMMARY . . . . .	v
Section	
1. INTRODUCTION. . . . .	1
2. THE ONE-DIMENSIONAL HEAT EQUATION . . . . .	2
3. THE EQUATION $u_t = uu_x$ . . . . .	3
REFERENCES. . . . .	5

SOME QUESTIONS CONCERNING DIFFERENCE APPROXIMATIONS  
TO PARTIAL DIFFERENTIAL EQUATIONS

1. INTRODUCTION

The study of finite-difference approximations to partial differential equations is well advanced, and the increasing use and significance of digital computers guarantees that this study will continue to be actively pursued. One question, however, that does not seem to have been discussed to any extent is that of finding discrete approximations that manifestly exhibit the boundedness, positivity, and so on, of the original continuous operator.

Thus, for example, the recurrence relation

$$(1.1) \quad u(t + \Delta) = (1 - a\Delta)u(t), \quad u(0) = c,$$

yields the nonnegativity we associate with the solution of

$$(1.2) \quad u' = -au, \quad u(0) = c,$$

provided that  $a\Delta < 1$ . On the other hand, the relation

$$(1.3) \quad u(x, t + \Delta^2) = \frac{u(x + \Delta\sqrt{2}, t) + u(x - \Delta\sqrt{2}, t)}{2},$$

$t = 0, \Delta^2, \dots$ , shows very clearly the nonnegativity associated with the solution of

$$(1.4) \quad u_t = u_{xx}, \quad u(x, 0) = g(x).$$

Approximations of this type are very useful computationally because of their stability properties. We shall discuss



briefly below the problem of obtaining approximations of arbitrarily high degree with the same property, and the related problem for

$$(1.5) \quad u_t = uu_x + g(u).$$

For previous work and computational results, see [1].

## 2. THE ONE-DIMENSIONAL HEAT EQUATION

To obtain higher-order approximations possessing the property that nonnegativity is preserved, we can proceed as follows. Write

$$(2.1) \quad u(x, t + \Delta^2) = \sum_{i=1}^N w_i (u(x + r_i \Delta, t) + u(x - r_i \Delta, t)).$$

Using the relations  $u_t = u_{xx}$ ,  $u_{tt} = u_{xxxx}$ , and so on, we obtain a set of moment relations

$$(2.2) \quad 1 = \sum_1 w_i,$$

$$\frac{1}{2} = \sum_1 w_i r_i^2,$$

and so on. The question of existence of real  $r_i$  and positive  $w_i$  yielding an error which is  $O(\Delta^{2k})$  for a given  $k$  and  $N$  can then be determined by invoking classical moment theory.

More interesting would be a general result to the effect that any linear partial differential equation

with constant coefficients can be approximated by difference relations of arbitrarily high degree which preserve the nonnegativity and boundedness properties of the original equation. Let us further note that a nonlinear approximation, e.g., one arising from a branching process, may be superior in that it requires fewer terms to obtain an equivalent approximation.

3. THE EQUATION  $u_t = uu_x$

Consider now the equation

$$(3.1) \quad u_t = uu_x, \quad u(x,0) = g(x),$$

which is often used (see [1]) as a simple model of an equation generating a shock wave. Since it possesses an explicit analytic solution, it is useful in testing numerical integration techniques. An approximation to terms in  $O(\Delta^2)$  is obtained using the difference relation

$$(3.2) \quad u(x, t + \Delta) = u(x + u(x, t)\Delta, t),$$

$$t = 0, \Delta, \dots$$

An approximation to terms in  $O(\Delta^2)$  is obtained from

$$(3.3) \quad u(x, t + \Delta) = u(x + u(x + u(x, t)\Delta, t)\Delta, t).$$

One might suspect that arbitrarily accurate approximations could be obtained by continuing in this fashion, but it seems difficult to establish this.

Similarly, the equation

$$(3.4) \quad u_t = uu_x + g(u)$$

has the approximation

$$(3.5) \quad u(x, t + \Delta) = u(x + u(x, t)\Delta, t) + g(u)\Delta,$$

valid to  $O(\Delta^2)$ , and

$$(3.6) \quad u(x, t + \Delta) = u(x + u(x + u(x, t)\Delta, t)\Delta + \frac{g(u)\Delta^2}{2}, t) \\ + g(u(x + u(x, t)\Delta)\Delta + \frac{g'(u)g(u)\Delta^2}{2},$$

valid to  $O(\Delta^3)$ .

REFERENCE

1. Bellman, R., I. Cherry, and G. M. Wing, "A Note on the Numerical Integration of a Class of Nonlinear Hyperbolic Equations," Quarterly of Applied Mathematics, Vol. 16, 1958, pp. 181-183.